

CLENPIQ™

(sodium picosulfate, magnesium oxide,
and anhydrous citric acid) Oral Solution

10 mg/3.5 g/12 g per 160 mL bottle

TAKING CLENPIQ

STEPS TO COMPLETE THE PREP

Day-Before-Dose Regimen

Afternoon/Early Evening Before

BETWEEN 4-6 PM



Drink 40 oz (5 cups) of clear liquids.*
Finish liquids over the next 5 hours.

Evening Before

5 HOURS BEFORE PROCEDURE



Drink at least 24 oz (3 cups) of clear liquids over the next 5 hours.
Finish liquids 2 hours before your colonoscopy
or as advised by your doctor.



Hydration is important and it's part of the prep. Make sure to hydrate
before you take the prep, while you're taking the prep, and after the prep.

Prep Assistant Use the prep assistant as a guide for completing your prep.

**ON THE DAY
BEFORE THE
COLONOSCOPY,**
stop eating all solid
food and dairy, and
start hydrating by
drinking
clear liquids.



Afternoon/Early Evening before your colonoscopy

4-6 PM

Drink one bottle of CLENPIQ
Drink CLENPIQ right from the bottle

Then, keep hydrating

Drink 5 cups (8 oz each) of clear liquid.
Check off the cups below as you go!



Start: _____:_____

Finish: _____:_____

Evening before your colonoscopy

5 HOURS BEFORE PROCEDURE

Drink the other bottle of CLENPIQ
Drink CLENPIQ right from the bottle

Then, keep hydrating

Drink at least 3 cups (8 oz each) of clear liquid.
Check off the cups below as you go!



OR MORE

Start: _____:_____

Finish: _____:_____

*After your first dose, if severe bloating, swelling, or stomach pain occurs, delay the second dose until the symptoms resolve.

Please see reverse page for additional information about CLENPIQ, the importance of a clear liquid diet, and hydration.

INDICATION

CLENPIQ is a prescription medicine used by adults to clean the colon before a colonoscopy. CLENPIQ cleans your colon by causing you to have diarrhea. Cleaning your colon helps your healthcare provider see the inside of your colon more clearly during your colonoscopy.

IMPORTANT SAFETY INFORMATION

- Do not take CLENPIQ if your healthcare provider has told you that you have serious kidney problems, a blockage in your intestine (bowel obstruction), an opening in the wall of your stomach or intestines (bowel perforation), a very dilated intestine (toxic megacolon), problems with the emptying of food and fluid from your stomach (gastric retention), or an allergy to any of the ingredients in CLENPIQ.
- CLENPIQ and other bowel preparations can cause serious side effects, including serious loss of body fluid (dehydration) and changes in blood salts (electrolytes) in your blood. These changes can cause abnormal heartbeats that may result in death, seizures (this can happen even if you have never had a seizure), or kidney problems. Your chance of having fluid loss and changes in blood salts with CLENPIQ is higher if you have heart problems, have kidney problems, or take water pills or non-steroidal anti-inflammatory drugs (NSAIDs).

See complete Important Safety Information on the back.