

2-Day Split-Dosing

If you're starting PLENVU the **evening before** your colonoscopy, **Here's How to Prep**

Date of colonoscopy://		Location of colonoscopy:	
Time of colonoscopy: AM/PM Nothing to drink after: AM/PM			
Evening		Morning	
PLENVU Dose 1:		PLENVU Dose 2:	
6 pm the evening before procedure GasX 1-2 drops/tablet w/ consuming prep		4-5 hours before arrival at facility GasX 1-2 drops/tablet w/ consuming prep	
EVENING 🗡 DOSE 1	MORNING XXXX DOSE 2		Stay hydrated! It's important that you drink
16 oz	16 oz 16 oz	READY FOR COLONOSCOPY	clear liquids before, during, and after your prep

Note: actual solution is clear.

MANGO FLAVOR



- 1. Use the mixing container to mix the contents of the Dose 1 pouch with at least 16 ounces of water by shaking or using a spoon until it's completely dissolved. This may take up to 2 to 3 minutes. Take your time—slowly finish the dose within 30 minutes.
- 2. Refill the container with at least 16 ounces of clear liquid. Again, take your time and slowly finish all of it within 30 minutes. Continue to drink additional clear liquids throughout the evening.

FRUIT PUNCH FLAVOR

Take Dose 2:

- Use the mixing container to mix the contents of Dose 2 (Pouch A and Pouch B) with at least 16 ounces of water by shaking or using a spoon until it's completely dissolved. This may take up to 2 to 3 minutes. Take your time—slowly finish the dose within 30 minutes.
- Refill the container with at least 16 ounces of clear liquid. Again, take your time and slowly finish all of it within 30 minutes. Continue
 to drink additional clear liquids, but be sure to stop drinking liquids at least 2 hours before your colonoscopy or as recommended by
 your healthcare provider.

For full instructions, refer to the Medication Guide and Instructions for Use in the Prescribing Information.

HELPFUL TIP: PLENVU can be refrigerated. Use within 6 hours after it's mixed with water

INDICATION

PLENVU[®] (polyethylene glycol 3350, sodium ascorbate, sodium sulfate, ascorbic acid, sodium chloride, and potassium chloride for oral solution) is a prescription medication used by adults to clean the colon before a colonoscopy.

IMPORTANT SAFETY INFORMATION

• Do not take PLENVU if you have a blockage in your intestine (bowel obstruction), an opening in the wall of your stomach or intestine

(bowel perforation), problems with food or fluid emptying from your stomach (gastric retention), a problem with food moving too slowly through your intestines (ileus), a very dilated large intestine, or an allergy to any of the ingredients in PLENVU.

Please see additional Important Safety Information on back and accompanying full Prescribing Information, including Medication Guide and Instructions for Use.