

## **GoLytely Prep Instructions**

## Day Before Procedure:

- □ 1 DAY PREP- Clear Liquid Diet (If you cannot see through it, DO NOT have it)
- □ GoLytely Prep

•Fill the bottle with lukewarm water until you reach the top fill line on the bottle.

•Recap the bottle and shake well to dissolve the powder completely. Once mixed, solution should be clear and colorless.

•Keep solution refrigerated until you are ready to drink. The reconstituted solution should be used within 48 hours.

## Consuming the solution- Start at 4:00 pm

- Drink 8 oz of solution every 10 minutes. The entire gallon must be consumed within 4 hours. Tip: Rapid drinking of each portion is better than drinking small amounts continuously.
- □ The first bowel movement should occur approximately within 1-2 hours after you start drinking the solution.
- Continue to drink the GoLytely solution until your bowel movements consist of watery stool that is clear and free of solid matter. This usually happens after you've drank at least 3 liters.
- ☐ You will have loose bowel movements for about 1 to 2 hours after you finish drinking the solution.
- □ Make sure you have drunk the entire contents of the bottle

## <u>At 10:00 pm</u>

- □ If stool isNOT COMPLETELY LIQUID →Take two (2) tablets of Dulcolax
- Stool should look similar to urine *no drinking after midnight*

*If you are taking essential meds like BP and seizure meds you may take with sip or water in am Any questions about the instructions or need further assistance? contact our office*